



Solving Quadratic Diophantine Equation for Integral Powers of 37

M. Vasuki¹, Dr. R. Sivaraman²

¹Assistant Professor, Department of Mathematics Srinivasan College of Arts and Science (Affiliated to Bharathidasan University) Perambalur, Tamil Nadu, India

²Associate Professor, Post Graduate & Research Department of Mathematics Dwaraka Doss Goverdhan Doss Vaishnav College, Chennai, India

ARTICLE INFO	ABSTRACT
Published Online: 31 January 2024	Diophantine Equations named after ancient Greek mathematician Diophantus, plays a vital role not only in number theory but also in several branches of science. In this paper, we have solved an quadratic Diophantine equations where the right hand side are positive integral powers of 37 and provide its integer solutions. The method adopted to solve the given equation is using the concept of polar form of a particular complex number. This concept can be generalized for solving similar equations.
Corresponding Author: Dr. R. Sivaraman	
KEYWORDS: Quadratic Diophantine Equation, Polar Form, Euler's Formula, Positive Integer Solutions.	

1. INTRODUCTION

Diophantine Equations were equations whose solutions must be in integers. Since the solutions are integers and most often positive integers, such equations have more practical applications compared to other equations in mathematics. In this paper, we will solve one of the quadratic Diophantine equations involving positive integral powers of 37 in a novel way and present its complete solution in a compact form.

2. QUADRATIC DIOPHANTINE EQUATION

the polar form of $(5 + i2\sqrt{3})^n$ $5 + i2\sqrt{3} = r(\cos \theta + i \sin \theta) \Rightarrow r \cos \theta = 5, r \sin \theta = 2\sqrt{3}$

From this, we obtain $r^2 = 25 + 12 = 37 \Rightarrow r = \sqrt{37}, \theta = \tan^{-1}\left(\frac{2\sqrt{3}}{5}\right)$ (2)

Hence the polar form of $(5 + i2\sqrt{3})^n$ is given by

$$(5 + i2\sqrt{3})^n = 37^{n/2} e^{i n \tan^{-1}\left(\frac{2\sqrt{3}}{5}\right)} \quad (3)$$

Now using Euler's Formula in (3), we obtain

$$(5 + i2\sqrt{3})^n = 37^{n/2} \left[\cos\left(n \tan^{-1}\left(\frac{2\sqrt{3}}{5}\right)\right) + i \sin\left(n \tan^{-1}\left(\frac{2\sqrt{3}}{5}\right)\right) \right] \quad (4)$$

If we now assume $y + i2\sqrt{3}x = (5 + i2\sqrt{3})^n$ (5) then $y - i2\sqrt{3}x = (5 - i2\sqrt{3})^n$ (6)

In this paper, we will provide methods to solve the quadratic Diophantine equation $12x^2 + y^2 = 37^n$ (1), where x, y are positive integers. We will try to obtain a general solution of (1) in closed form. For doing this, we will make use of a particular complex number and a wonderful method proposed by the greatest mathematician of all times, Leonhard Euler.

3. SOLUTIONS TO THE EQUATION

We will try to obtain all positive integer solutions (x, y) satisfying (1) for any given natural number n . Now, we will try to determine

Now multiplying (5) and (6), we get

$$(y + i2\sqrt{3}x) \times (y - i2\sqrt{3}x) = (5 + i2\sqrt{3})^n \times (5 - i2\sqrt{3})^n$$

Simplifying, we obtain $12x^2 + y^2 = 37^n$ which is (1), the original problem which we have considered. Thus the solutions to (1) are given by equating real and imaginary parts of (5). Now using (4) in (5), and for $n \geq 1$ we get

$$2\sqrt{3}x = 37^{n/2} \sin\left(n \tan^{-1}\left(\frac{2\sqrt{3}}{5}\right)\right) \Rightarrow x = \frac{37^{n/2}}{2\sqrt{3}} \sin\left(n \tan^{-1}\left(\frac{2\sqrt{3}}{5}\right)\right) \quad (7)$$

$$y = 37^{n/2} \cos\left(n \tan^{-1}\left(\frac{2\sqrt{3}}{5}\right)\right) \Rightarrow y = 37^{n/2} \cos\left(n \tan^{-1}\left(\frac{2\sqrt{3}}{5}\right)\right) \quad (8)$$

Now from (7) and (8), if we consider $(|x|, |y|)$ then these pairs would provide all positive integer solutions to the given Quadratic Diophantine Equation $12x^2 + y^2 = 37^n$ for any natural number n .

4. CONCLUSION

Considering a quadratic Diophantine equation $12x^2 + y^2 = 37^n$, we have used a novel method to solve it completely in this paper. In particular, equations (7) and (8) provide all required positive integer solutions to the given equation. Further, by considering the polar form of a particular complex number, I have obtained nice closed expressions for the given equations.

In fact, from (7) and (8), we notice that for $n \geq 1$, all positive integer solutions to $12x^2 + y^2 = 37^n$ are given by

$$x = \frac{37^{n/2}}{2\sqrt{3}} \left| \sin\left(n \tan^{-1}\left(\frac{2\sqrt{3}}{5}\right)\right) \right|, y = 37^{n/2} \left| \cos\left(n \tan^{-1}\left(\frac{2\sqrt{3}}{5}\right)\right) \right| \quad (9)$$

Thus, for $n = 1, 2, 3, 4, 5, 6, 7, 8, \dots$ all positive integer solutions (x, y) to $12x^2 + y^2 = 37^n$ are given respectively by (1,5); (10,13); (63,55); (260,1031); (269,8275); (6930,44603); (79253,139855); (536120,251761); \dots

Thus the values of x and y from expression (9) provides all possible positive integer solutions to the given quadratic Diophantine equation $12x^2 + y^2 = 37^n$. We can adopt similar methods to solve other types of quadratic Diophantine equations using polar forms of suitable complex numbers.

REFERENCES

1. Andreescu, T., D. Andrica, and I. Cucurezeanu, An introduction to Diophantine equations: A problem-based approach, Birkhäuser Verlag, New York, 2010.
2. Andrews, G. E. 1971, Number theory, W. B. Saunders Co., Philadelphia, Pa.-London- Toronto, Ont.
3. R. Sivaraman, J. Suganthi, A. Dinesh Kumar, P.N. Vijayakumar, R. Sengothai, On Solving an Amusing Puzzle, Specialusis Ugdymas/Special Education, Vol 1, No. 43, 2022, 643 – 647.
4. R. Sivaraman, R. Sengothai, P.N. Vijayakumar, Novel Method of Solving Linear Diophantine Equation with Three Variables, Stochastic Modeling and Applications, Vol. 26, No. 3, Special Issue – Part 4, 2022, 284 – 286.
5. R. Sivaraman, On Solving Special Type of Linear Diophantine Equation, International Journal of Natural Sciences, Volume 12, Issue 70, 38217 – 38219, 2022.
6. R. Sivaraman, Summing Through Integrals, Science Technology and Development, Volume IX, Issue IV, April 2020, pp. 267 – 272.
7. R. Sivaraman, Bernoulli Polynomials and Ramanujan Summation, Proceedings of First International Conference on Mathematical Modeling and Computational Science, Advances in Intelligent Systems and Computing, Vol. 1292, Springer Nature, 2021, pp. 475 – 484.
8. R. Sengothai, R. Sivaraman, Solving Diophantine Equations using Bronze Ratio, Journal of Algebraic Statistics, Volume 13, No. 3, 2022, 812 – 814.
9. P.N.Vijayakumar, R. Sivaraman, On Solving Euler’s Quadratic Diophantine Equation, Journal of Algebraic Statistics, Volume 13, No. 3, 2022, 815 – 817.
10. A. Dinesh Kumar, R. Sivaraman, Asymptotic Behavior of Limiting Ratios of Generalized Recurrence Relations, Journal
11. P Senthil Kumar, R Abirami, A Dinesh Kumar, Fuzzy model for the effect of rhIL6 Infusion on

“Solving Quadratic Diophantine Equation for Integral Powers of 37”

- Growth Hormone, International Conference on Advances in Applied Probability, Graph Theory and Fuzzy Mathematics, Vol. 252, 2014, pp. 246
12. P Senthil Kumar, A Dinesh Kumar, M Vasuki, Stochastic model to find the effect of gallbladder contraction result using uniform distribution, Arya Bhatta Journal of Mathematics and Informatics, Vol 6, No. 2, 2014, pp. 323 – 328
 13. R. Sivaraman, Recognizing Ramanujan’s House Number Puzzle, German International Journal of Modern Science, 22, November 2021, pp. 25 – 27
 14. R. Sivaraman, Pythagorean Triples and Generalized Recursive Sequences, Mathematical Sciences International Research Journal, Vol. 10, No. 2, July 2021, pp. 1 – 5
 15. R. Sivaraman, Sum of powers of natural numbers, AUT AUT Research Journal, Volume XI, Issue IV, April 2020, pp. 353 – 359.